

Pediculosis Pubis

This is a PDF version of the following document:

Module 1: [STD Question Bank](#)

Lesson 20: [Pediculosis Pubis](#)

You can always find the most up-to-date version of this document at

<https://www.std.uw.edu/go/2021-guidelines/pediculosis-pubis/core-concept/all>.

**Table 1. 2021 STI Treatment Guidelines: Ectoparasitic Infections
Treatment of Pediculosis Pubis**

Recommended for Treatment of Pediculosis Pubis

Permethrin 1% cream rinse

applied to affected areas and washed off after 10 minutes

Recommended for Treatment of Pediculosis Pubis

Pyrethrin with piperonyl butoxide

applied to the affected area and washed off after 10 minutes

Alternative for Treatment of Pediculosis Pubis

Malathion 0.5% lotion

applied to affected areas and washed off after 8–12 hours

Alternative for Treatment of Pediculosis Pubis

Ivermectin

250 µg/kg body weight, orally, repeated in 7 to 14 days

Source: Workowski KA, Bachmann LH, Chan PA, et al. Sexually transmitted infections treatment guidelines, 2021. Ectoparasitic infections. MMWR Recomm Rep. 2021;70(No. RR-4):1-187. [[2021 STI Treatment Guidelines](#)]

